

## D-Dimer antibody [DD2]

Cat. No. GTX10052

宿主	Mouse
克隆	Monoclonal
同种型	IgG2b
实验应用	WB, ELISA, Immunoassay, Sandwich ELISA
种属反应	Human

## 实验应用

## 应用说明

\*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	Assay dependent
ELISA	Assay dependent
Immunoassay	Assay dependent
Sandwich ELISA	Assay dependent

**Note : Immunoassays for the quantitative determination of D-dimer and high molecular weight fibrin degradation products.**

**Capture : GTX10052, Detection : GTX05174 / GTX43438 or Capture : GTX10052, Detection : GTX10054**

**Due to the cross-reactivity of DD4 with fibrinogen, we strongly recommend to use it as the detection antibody. In a sandwich immunoassay, plasma must be diluted at least two-fold with 10 mM Tris-HCl, pH 7.5, 1 M NaCl, 0.1 % Tween 20 in order to avoid nonspecific binding. Each step in the assay should be followed by an incubation and wash: coating with the capture antibody, addition of the sample, and addition of the detection antibody.**

**Please notice that the detected antibody needs to be conjugated to HRP to function as the detection antibody when paired with the capture antibody.**

**Please contact us for custom HRP-conjugated antibody.**

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片

## 属性

形式	Liquid
存储溶液	PBS
保存剂	0.09% Sodium azide
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
浓度	Batch dependent (Please refer to the vial label for the specific concentration.)



For full product information, images and publications, please visit our [website](#).

---

偶联 Unconjugated

---

注意事项 仅供实验室使用。不适用于人类或动物的任何临床, 治疗或诊断用途。不适合动物或人类食用。

---



For full product information, images and publications, please visit our [website](#).