

D-Dimer antibody [DD4]

Cat. No. GTX10054

宿主	Mouse
克隆	Monoclonal
同种型	IgG2b
实验应用	WB, ELISA, Immunoassay, Sandwich ELISA
种属反应	Human

引用文献 (2)

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	Assay dependent
ELISA	Assay dependent
Immunoassay	Assay dependent
Sandwich ELISA	Assay dependent

Note : Immunoassays for the quantitative determination of D-dimer and high molecular weight fibrin degradation products.

Capture : GTX10052 ; Detection : GTX10054, Due to the cross-reactivity of DD4 with fibrinogen, we strongly recommend to use it as the detection antibody. In a sandwich immunoassay, plasma must be diluted at least two-fold with 10 mM Tris-HCl, pH 7.5, 1 M NaCl, 0.1 % Tween 20 in order to avoid nonspecific binding. Each step in the assay should be followed by an incubation and wash: coating with the capture antibody, addition of the sample, and addition of the detection antibody.

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片

属性

形式	Liquid
存储溶液	PBS
保存剂	0.09% Sodium azide
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. Store at 4°C.
浓度	Batch dependent (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
注意事项	仅供实验室使用。不适用于人类或动物的任何临床、治疗或诊断用途。不适合动物或人类食用。



For full product information, images and publications, please visit our [website](#).