

## AKAP10 antibody [C3], C-term

**Cat. No. GTX106430**

宿主	Rabbit
克隆	Polyclonal
同种型	IgG
实验应用	ICC/IF, IHC-P
种属反应	Human, Mouse, Rat

## 实验应用

## 应用说明

\*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片


## 属性

形式	Liquid
存储溶液	0.1M Tris, 0.1M Glycine, 10% Glycerol
保存剂	0.01% Thimerosal
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓度	0.35 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
RRID	AB_1949618
注意事项	仅供实验室使用。不适用于人类或动物的任何临床, 治疗或诊断用途。不适合动物或人类食用。




For full product information, images and publications, please visit our [website](#).

## 產品圖片

**GTX106430 IHC-P Image**

Immunohistochemical analysis of paraffin-embedded human OVCAR3, using AKAP10(GTX106430) antibody at 1:100 dilution.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min


**GTX106430 IHC-P Image**

AKAP10 antibody [C3], C-term detects AKAP10 protein at cytoplasm in rat liver by immunohistochemical analysis.


Sample: Paraffin-embedded rat liver.

AKAP10 antibody [C3], C-term (GTX106430) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min

**GTX106430 ICC/IF Image**

Immunofluorescence analysis of methanol-fixed HeLa, using AKAP10(GTX106430) antibody at 1:500 dilution.

**GTX106430 IHC-P Image**

AKAP10 antibody [C3], C-term detects AKAP10 protein at cytoplasm in mouse brain by immunohistochemical analysis.

Sample: Paraffin-embedded mouse brain.

AKAP10 antibody [C3], C-term (GTX106430) diluted at 1:100.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



For full product information, images and publications, please visit our [website](#).