

C20orf7 antibody [N3C3]

Cat. No. GTX119214

宿主	Rabbit
克隆	Polyclonal
同种型	IgG
实验应用	WB, ICC/IF, IHC-P
种属反应	Human

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	1:100-1:1000
IHC-P	1:100-1:1000

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片


属性

形式	Liquid
存储溶液	0.1M Tris, 0.1M Glycine, 20% Glycerol
保存剂	0.01% Thimerosal
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓度	0.84 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
RRID	AB_10724480
注意事项	仅供实验室使用。不适用于人类或动物的任何临床、治疗或诊断用途。不适合动物或人类食用。




For full product information, images and publications, please visit our [website](#).

產品圖片




GTX119214 ICC/IF Image

C20orf7 antibody [N3C3] detects C20orf7 protein at cytoplasm by immunofluorescent analysis.
 Sample: SK-N-SH cells were fixed in 2% paraformaldehyde/culture medium at 37°C for 30 min.
 Green: C20orf7 protein stained by C20orf7 antibody [N3C3] (GTX119214) diluted at 1:500.
 Blue: Hoechst 33342 staining.



GTX119214 WB Image

Sample (30 ug of whole cell lysate)
 A: IMR32
 10% SDS PAGE
 GTX119214 diluted at 1:2000



GTX119214 IHC-P Image

Immunohistochemical analysis of paraffin-embedded human gastric cancer, using C20orf7(GTX119214) antibody at 1:500 dilution.
 Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



For full product information, images and publications, please visit our [website](https://www.genetex.com).