

Histone H3R17me2 (Asymmetric Di-methyl Arg17) antibody

Cat. No. GTX124113

| | |
|------|----------------|
| 宿主 | Rabbit |
| 克隆 | Polyclonal |
| 同种型 | IgG |
| 实验应用 | WB, IHC-P, Dot |
| 种属反应 | Human, Mouse |

引用文献 (1)

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB | 1:500-1:3000 |
| IHC-P | 1:100-1:1000 |
| Dot | Assay dependent |

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片


IHC-Fr: 冰冻切片

属性

| | |
|------|--|
| 形式 | Liquid |
| 存储溶液 | PBS, 1% BSA, 20% Glycerol |
| 保存剂 | 0.01% Thimerosal |
| 存放说明 | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| 浓度 | 0.62 mg/ml (Please refer to the vial label for the specific concentration.) |
| 偶联 | Unconjugated |
| RRID | AB_2885596 |
| 注意事项 | 仅供实验室使用。不适用于人类或动物的任何临床, 治疗或诊断用途。不适合动物或人类食用。 |

For full product information, images and publications, please visit our [website](#).

產品圖片

**GTX124113 WB Image**

Histone3 (di-methyl Arg17) antibody detects HIST1H3A protein by Western blot analysis.

- A. 30 μ g 293T whole cell lysate/extract
- B. 30 μ g A431 whole cell lysate/extract
- C. 30 μ g HeLa whole cell lysate/extract
- D. 30 μ g HepG2 whole cell lysate/extract

15 % SDS-PAGE

Histone3 (di-methyl Arg17) antibody (GTX124113) dilution: 1:1000


**GTX124113 IHC-P Image**

Histone H3R17me2 (dimethyl Arg17) antibody detects Histone H3R17me2 (dimethyl Arg17) protein at nucleus on mouse muscle by immunohistochemical analysis.

Sample: Paraffin-embedded mouse muscle.

Histone H3R17me2 (dimethyl Arg17) antibody (GTX124113) dilution: 1:500.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min


**GTX124113 IHC-P Image**

Histone3 (di-methyl Arg17) antibody detects HIST1H3A protein at nucleus on AGS xenograft by immunohistochemical analysis.

Sample: Paraffin-embedded AGS xenograft.

Histone3 (di-methyl Arg17) antibody (GTX124113) dilution: 1:500.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min

**GTX124113 IHC-P Image**

Histone H3R17me2 (dimethyl Arg17) antibody detects Histone H3R17me2 (dimethyl Arg17) protein at nucleus on mouse middle brain by immunohistochemical analysis.

Sample: Paraffin-embedded mouse middle brain.

Histone H3R17me2 (dimethyl Arg17) antibody (GTX124113) dilution: 1:500.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



For full product information, images and publications, please visit our [website](#).