

## Influenza A virus H3N2 HA (Hemagglutinin) antibody

Cat. No. GTX127363

宿主	Rabbit
克隆	Polyclonal
同种型	IgG
实验应用	WB, ICC/IF, IHC-P
种属反应	Influenza A virus

引用文献 (10)

## 实验应用

## 应用说明

\*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	1:500-1:3000
ICC/IF	Assay dependent
IHC-P	1:50-1:1000

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片


IHC-Fr: 冰冻切片

## 属性

形式	Liquid
存储溶液	PBS, 1% BSA, 20% Glycerol
保存剂	0.025% ProClin 300
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓度	0.23 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
RRID	AB_2885648
注意事项	仅供实验室使用。不适用于人类或动物的任何临床, 治疗或诊断用途。不适合动物或人类食用。


For full product information, images and publications, please visit our [website](#).

產品圖片



**GTX127363 WB Image**


Non-transfected (-) and transfected (+) 293T whole cell extracts (30 µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with Influenza A virus H3N2 HA (Hemagglutinin) antibody (GTX127363) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



**GTX127363 IHC-P Image**

Influenza A Virus H3N2 Hemagglutinin (HA) antibody detects Influenza A Virus H3N2 Hemagglutinin (HA) protein on mouse nasal tissue (infect with Influenza Virus A) by immunohistochemical analysis. Sample: Paraffin-embedded mouse nasal tissue (infect with Influenza Virus A). Influenza A Virus H3N2 Hemagglutinin (HA) antibody (GTX127363) dilution: 1:50.

Antigen Retrieval: Trilogy™ (EDTA based, pH 8.0) buffer, 15min



**GTX127363 WB Image**


Influenza A Virus H3N2 Hemagglutinin (HA) antibody detects HA (H3N2) protein by western blot analysis.

- A. MDCK cell lysates infected with 4 HAU/ml of A/duck/Alberta/35/1976-like, H1N1
- B. MDCK cell lysates infected with 4 HAU/ml of A/duck/Ukraine/1963-like, H3N8
- C. MDCK cell lysates infected with 4 HAU/ml of A/mallard/Miyagi/53/1976-like, H5N3
- D. MDCK cell lysates infected with 4 HAU/ml of A/duck/Taiwan/33/1993-like, H7N7
- E. MDCK cell lysates infected with 4 HAU/ml of A/turkey/Wisconsin/1/1966-like, H9N2
- F. MDCK cell lysates infected with 4 HAU/ml of A/PR/8/34, H1N1
- G. MDCK cell lysates infected with 4 HAU/ml of A/Taiwan/126/2009, pdmH1N1
- H. MDCK cell lysates infected with 4 HAU/ml of A/Taiwan/3446/2002, H3N2

12% SDS-PAGE

Influenza A Virus H3N2 Hemagglutinin (HA) antibody (GTX127363) dilution: 1:1000

The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



**GTX127363 WB Image**

Recombinant H3N2 HA protein were separated by 10% SDS-PAGE, and the membrane was blotted with Influenza A virus H3N2 HA (Hemagglutinin) antibody (GTX127363) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody.



For full product information, images and publications, please visit our [website](#).