

JNK (phospho Thr183/Tyr185) antibody

Cat. No. GTX133934

宿主	Rabbit
克隆	Polyclonal
同种型	IgG
实验应用	WB, IHC-P
种属反应	Human, Mouse, Drosophila

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	1:1000
IHC-P	Assay dependent

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片


属性

形式	Liquid
存储溶液	PBS, 1% BSA, 20% Glycerol
保存剂	0.025% ProClin 300
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓度	1.09 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
RRID	AB_2887139
注意事项	仅供实验室使用。不适用于人类或动物的任何临床, 治疗或诊断用途。不适合动物或人类食用。




For full product information, images and publications, please visit our [website](#).


產品圖片

**GTX133934 WB Image**

Untreated (-) and treated (+) 293T whole cell extracts (30 μg) were separated by 10% SDS-PAGE, and the membrane was blotted with JNK (phospho Thr183/Tyr185) antibody (GTX133934) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.

**GTX133934 WB Image**

Drosophila tissue extract (50 μg) was separated by 10% SDS-PAGE, and the membrane was blotted with JNK (phospho Thr183/Tyr185) antibody (GTX133934) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.


**GTX133934 IHC-P Image**

JNK (phospho Thr183/Tyr185) antibody detects JNK (phospho Thr183/Tyr185) protein at nucleus by immunohistochemical analysis.

Sample: Paraffin-embedded human breast carcinoma.

JNK (phospho Thr183/Tyr185) stained by JNK (phospho Thr183/Tyr185) antibody (GTX133934) diluted at 1:500.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min

**GTX133934 WB Image**

Untreated (-) and treated (+) Raw264.7 whole cell extracts (30 μg) were separated by 10% SDS-PAGE, and the membrane was blotted with JNK (phospho Thr183/Tyr185) antibody (GTX133934) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.



For full product information, images and publications, please visit our [website](#).