

PCNA antibody [PC10]

Cat. No. GTX20029

宿主	Mouse
克隆	Monoclonal
同种型	IgG2a
实验应用	WB, ICC/IF, IHC-P, IHC-Fr, FCM, IP, IHC
种属反应	Human, Mouse, Rat, Rabbit, Fish

引用文献 (27)

实验应用


应用说明

Flow Cyt: Use at an assay dependent dilution. IHC-P: 1/1600. Perform heat mediated antigen retrieval before commencing with IHC staining protocol
IHC-Fr: Use at an assay dependent dilution. Fix in 4% buffered formalin at room temperature for 2 min then fix in ethanol for 10 minutes. IP: Use at an assay dependent dilution WB: 1:5,000, Detects a band of approximately 30 kDa (predicted molecular weight: 29 kDa). Optimal dilutions/concentrations should be determined by the researcher.

属性

形式	Liquid
存储溶液	PBS
保存剂	No preservatives
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓度	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
RRID	AB_369854
注意事项	仅供实验室使用。不适用于人类或动物的任何临床，治疗或诊断用途。不适合动物或人类食用。

产品图片




GTX20029 ICC/IF Image


Human Tonsil stained with Anti-CD3, T Cell at a 1:150 dilution (red) and Monoclonal Anti-Proliferating Cell Nuclear Antigen (GTX20029) at a 1:500 dilution (green). Counterstained with DAPI.




For full product information, images and publications, please visit our [website](#).


GTX20029 WB Image

Western blot analysis of HeLa Nuclear Extract probed with Mouse anti Human PCNA (GTX20029)


GTX20029 FCM Image

Staining of KM-H2 cells with MOUSE ANTI PCNA:FITC (GTX75144) following permeabilisation


GTX20029 IHC-P Image

The data was published in the journal J Clin Med in 2019. [PMID: 30871022](https://pubmed.ncbi.nlm.nih.gov/30871022/)



For full product information, images and publications, please visit our [website](#).