

PDZK3 antibody, N-term

Cat. No. GTX26037

宿主	Goat
克隆	Polyclonal
同种型	IgG
实验应用	ICC/IF, IHC-P
种属反应	Human

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
ICC/IF	Assay dependent
IHC-P	Assay dependent

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片


属性

形式	Liquid
存储溶液	TBS, 0.5% BSA
保存剂	0.02% Sodium azide
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓度	0.5 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
注意事项	仅供实验室使用。不适用于人类或动物的任何临床、治疗或诊断用途。不适合动物或人类食用。



For full product information, images and publications, please visit our [website](#).


產品图片

**GTX26037 IHC-P Image**

IHC-P analysis of human tonsil tissue using GTX26037 PDZK3 antibody, N-term.

Antigen retrieval : Heat induced antigen retrieval with citrate buffer pH 6

Dilution : 6µg/ml

**GTX26037 ICC/IF Image**

ICC/IF analysis of paraformaldehyde fixed U2OS cells using GTX26037 PDZK3 antibody, N-term.

Green : Primary antibody

Blue : DAPI

Negative control: Unimmunized goat IgG

Dilution : 10µg/ml

Green : Primary antibody

Blue : DAPI

Negative control: Unimmunized goat IgG

Dilution : 10µg/ml



For full product information, images and publications, please visit our [website](#).