

BCIP/NBT Substrate

Cat. No. GTX30943

实验应用 WB, ICC/IF, IHC, ISH

Package
50 ml

产品说明

摘要 BCIP/NBT Substrate

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
WB	Assay dependent
ICC/IF	Assay dependent
IHC	Assay dependent
ISH	Assay dependent

- Note :**
1. After the alkaline phosphatase reaction, wash slide with high pH buffer (~Tris/HCl 9.5).
 2. Add few drops of BCIP/NBT reagent, incubate at RT or 37°C for 5-15 minutes.
 3. Wash 2-5 times with buffer, followed by rinsing with distilled/deionized water.
 4. Add counterstain compatible with this chromogen (nuclear fast red)
 5. Wash with tap water, followed by rinsing with distilled/deionized water.
 6. Mount with aqueous mounting medium.
- Phosphate buffer is a strong inhibitor of alkaline phosphatase and should not be used.

以下为常规应用缩写的中文注解

WB: 免疫印迹
 ICC/IF: 细胞染色
 IHC-P: 石蜡切片
 IHC-Fr: 冰冻切片

属性

形式 Liquid

存放说明 Store at 2-8°C.

注意事项 仅供实验室使用。不适用于人类或动物的任何临床, 治疗或诊断用途。不适合动物或人类食用。



For full product information, images and publications, please visit our [website](#).