

## CD62P antibody [AK-6]

Cat. No. GTX31222

|      |                                |
|------|--------------------------------|
| 宿主   | Mouse                          |
| 克隆   | Monoclonal                     |
| 同种型  | IgG1                           |
| 实验应用 | WB, ICC/IF, IHC-P, IHC-Fr, FCM |
| 种属反应 | Human, Rhesus Monkey           |

## 实验应用

## 应用说明

\*最佳稀释倍数与浓度应由研究人员确认

| Suggested dilution | Recommended dilution |
|--------------------|----------------------|
| WB                 | Assay dependent      |
| ICC/IF             | Assay dependent      |
| IHC-P              | 1/500-1/2,000        |
| IHC-Fr             | 1/1,000-1/5,000      |
| FCM                | 1/10-1/100           |

**Note : This product may enhance using Sodium Citrate buffer pH6.0**  
**Use 10 $\mu$ l of the suggested working dilution to label 10<sup>6</sup> cells in 100 $\mu$ l.**

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片


## 属性

|      |  |
|------|--|
| 形式   | Liquid   |
| 存储溶液 | PBS  |
| 保存剂  | 0.09% Sodium azide   |
| 存放说明 | Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles. |
| 浓度   | 1.0 mg/ml (Please refer to the vial label for the specific concentration.)   |
| 偶联   | Unconjugated   |
| 注意事项 | 仅供实验室使用。不适用于人类或动物的任何临床，治疗或诊断用途。不适合动物或人类食用。   |




For full product information, images and publications, please visit our [website](#).

產品圖片




**GTX31222 FCM Image**

FACS analysis of human peripheral blood platelets using GTX31222 CD62P antibody [AK-6].



**GTX31222 IHC-P Image**

IHC-P analysis of human tonsil tissue using GTX31222 CD62P antibody [AK-6].



**GTX31222 FCM Image**

FACS analysis of thrombin activated human platelets using GTX31222 CD62P antibody [AK-6].



For full product information, images and publications, please visit our [website](#).