

CD4 antibody [OX-35]

Cat. No. GTX44519

宿主	Mouse
克隆	Monoclonal
同种型	IgG2a
实验应用	ICC/IF, IHC-Fr, FCM, IP
种属反应	Rat

实验应用

应用说明

*最佳稀释倍数与浓度应由研究人员确认

Suggested dilution	Recommended dilution
ICC/IF	Assay dependent
IHC-Fr	Assay dependent
FCM	1/10-1/50
IP	Assay dependent

Note : Use 10 μ l of the suggested working dilution to label 10⁶ cells in 100 μ l.

以下为常规应用缩写的中文注解

WB: 免疫印迹

ICC/IF: 细胞染色

IHC-P: 石蜡切片

IHC-Fr: 冰冻切片


属性

形式	Liquid
存储溶液	PBS
保存剂	0.09% Sodium azide
存放说明	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
浓度	1.0 mg/ml (Please refer to the vial label for the specific concentration.)
偶联	Unconjugated
注意事项	仅供实验室使用。不适用于人类或动物的任何临床, 治疗或诊断用途。不适合动物或人类食用。




For full product information, images and publications, please visit our [website](#).


產品圖片

**GTX44519 IHC-Fr Image**


IHC-Fr analysis of rat lymph node tissue using GTX44519 CD4 (Domain 2) antibody [OX-35].

**GTX44519 IHC-Fr Image**

IHC-Fr analysis of rat lymph node tissue using GTX44519 CD4 (Domain 2) antibody [OX-35].

**GTX44519 IHC-Fr Image**

IHC-Fr staining of rat lymph node using Mouse anti Rat CD4 antibody (GTX44519), red in A and Mouse anti Rat CD8, green in B. C is merged image with nuclear counterstaining with DAPI in blue. High power

**GTX44519 IHC-Fr Image**

IHC-Fr staining of rat lymph node using Mouse anti Rat CD4 antibody (GTX44519), red in A and Mouse anti Rat CD8, green in B. C is merged image with nuclear counterstaining with DAPI in blue. Low power

**GTX44519 IHC-Fr Image**

IHC-Fr analysis of rat lymph node tissue using GTX44519 CD4 (Domain 2) antibody [OX-35].



For full product information, images and publications, please visit our [website](#).